World Suicide Prevention Day

September 10th 2018

Working Together to Prevent Suicide

Facts and Figures





Over 800,000 people die by suicide annually, representing 1 person every 40 seconds







 Suicide is the 15th leading cause of death globally, account for 1.4% of all deaths and

• The global suicide rate is **11.4** per 100 000 population

15.0/100 000 for males

8.0/100 000 for females







September 10, 2018



 Suicide is the leading cause of death in people aged 15-24 in many European countries

 Globally suicide rates among this age group are higher in males than females

 Self-harm largely occurs among older adolescents, and globally is the 2nd leading cause of death for older adolescent girls





• In 2012, 76% of global suicide occurred in low- and middle-income countries 39% of which occurred in the South-East Asia Region





September 10, 2018



• In 25 countries (within WHO member states) suicide is currently still criminalized

• In an additional 20 countries suicide attempters may be punished with jail sentences, according to Sharia law





September 10, 2018



 Suicide is the result of a convergence of risk factors including but not limited to genetic, psychological, social and cultural risk factors, sometimes combined with experiences of trauma and loss





September 10, 2018



• Depression is the most common psychiatric disorder in people who die by suicide

• 50% of individuals in high income countries who die by suicide have major depressive disorder at their time of death





September 10, 2018

 For every 1 suicide 25 people make a suicide attempt





September 10, 2018



 135 people are affected by each suicide death

• This equates to 108 million people bereaved by suicide worldwide every year







Relatives and close friends of people who die by suicide are a high-risk group for suicide, due to:

- The psychological trauma of a suicide loss
- Potential shared familial and environmental risk
- Suicide contagion through the process of social modelling, and
- The burden of stigma associated with this loss





September 10, 2018

Suicide prevention strategies aim to prevent suicide among targeted high-risk groups but also at a universal level

Effective suicide prevention strategies need to incorporate public health policy strategies and healthcare strategies, incorporating measure with the strongest evidence of efficacy such as:

- Restriction of access to lethal means
- Treatment of depression
- Ensuring chain of care, and
- School-based universal prevention







September 10, 2018

Suicide prevention programmes face ongoing challenges including:

- Insufficient resources
- Ineffective coordination
- Limited access to surveillance data on suicide and self-harm
- Lack of enforced guidelines
- Lack of independent and systematic evaluation



September 10, 2018



 Since the 2014 WHO Global Report Preventing Suicide, progress in suicide prevention is ongoing, particularly in countries where no suicide initiatives were active, such as Guyana, Suriname and Bhutan



 Many countries have initiated their 2nd national suicide prevention programme, including:
 Scotland, England, Ireland and the United States





I A
S P
International
Association
for
Suicide
Prevention

 This World Suicide Prevention Day and everyday take a minute to reach out to someone - a complete stranger, close family member or friend - this can change a life





International Association for Suicide Prevention

September 10, 2018

The importance of community

 We are all part of a community, linked to family, friends, work colleagues, neighbours or teams.

- Sometimes we become isolated and disconnected from our communities.
- We have the responsibility to support and empower those who become vulnerable in our communities.



I A
S P
International
Association
for
Suicide
Prevention

September 10, 2018

Reaching out to those in the community

- If you are worried about someone in your community reach out and ask them "are you okay?"
- By simply checking in with them and offering nonjudgemental support you can make a difference.







September 10, 2018

Empowering our community

• Encourage those in distress to tell their own story in their way and at their pace.

 By engaging in active listening and reaching out to those who are vulnerable in the community together we can build resilient and strong

communities





September 10, 2018

No one has all of the answers

- People are often reluctant to intervene when they are worried for others.
- They may not feel knowledgeable enough or may not feel like the right person.
- It is important to know that people in distress are often not looking for specific advice, but merely to be listened to with compassion and empathy.





International Association for Suicide Prevention

September 10, 2018

Learn from people who reach out

- People with lived experience of suicidal behaviour or suicide bereavement have a lot to teach us.
- By listening to these often inspiring individuals we are learning and they may become empowered to seek help.



To find out more visit our WSPD webpage at: https://iasp.info/wspd2018/

Working Together to Prevent Suicide

References:

WHO, 2014; Mishara & Weisstub 2016; Arsenault-Lapierre et al., 2004; Cavanagh et al., 2003; Pitman et al., 2016; Zalsman et al., 2017